

ON QUANTUM INTEGRABILITY AND THE LEFSCHETZ NUMBER

Antti J. Niemi*

*Department of Theoretical Physics, Uppsala University
P.O. Box 108, S-75108, Uppsala, Sweden*

and

Kaupo Palo**

*Department of Theoretical Physics, Uppsala University
P.O. Box 108, S-75108, Uppsala, Sweden*

and

*Institute of Physics, Estonian Academy of Sciences
142 Riia St., 202 400 Tartu, Estonia*

Certain phase space path integrals can be evaluated exactly using equivariant cohomology and localization in the canonical loop space. Here we extend this to a general class of models. We consider hamiltonians which are *a priori* arbitrary functions of the Cartan subalgebra generators of a Lie group which is defined on the phase space. We evaluate the corresponding path integral and find that it is closely related to the infinitesimal Lefschetz number of a Dirac operator on the phase space. Our results indicate that equivariant characteristic classes could provide a natural geometric framework for understanding quantum integrability.

* E-mail: NIEMI@TETHIS.TEORFYS.UU.SE

** E-mail: PALO@RHEA.TEORFYS.UU.SE

There are several different approaches to quantum integrability [1,2]. Here we shall investigate a geometric approach which is based on phase space path integrals and loop space equivariant cohomology [3-13]. We show that the formalism is applicable to a general class of integrable models with hamiltonians that are functionals of canonical Lie algebra generators. Our result extends those derived previously in [12,13]. In particular, we find that for the class of models we consider the path integral can be evaluated exactly using localization techniques [3-13]. The final result is an ordinary integral over the phase space, and can be represented as a functional of an equivariant characteristic class [7]. We also find that the path integral can be related to the path integral representation of a Dirac operator which is defined on the phase space of the theory. In particular, the path integral of our integrable models is closely related to the infinitesimal Lefschetz number which is associated to this Dirac operator. Our results suggest that equivariant cohomology and equivariant characteristic classes seem to provide a natural framework for a geometric formulation of quantum integrability.

We consider a hamiltonian H on a phase space Γ with $2n$ coordinates z^a and Poisson bracket

$$\{z^a, z^b\}_\omega = \omega^{ab}(z) \quad (1)$$

Here the inverse matrix ω_{ab} determines the symplectic two-form on the phase space. We shall assume that on Γ there is a canonical realization of a Lie algebra by functions $G_u(z)$,

$$\{G_u, G_v\}_\omega = \omega^{ab} \partial_a G_u \partial_b G_v = f_{uv}{}^w G_w \quad (2)$$

and we shall assume that the hamiltonian H is a functional of these generators,

$$H(z) = H[G_u(z)] \quad (3)$$

An integrable hamiltonian H can be naturally viewed as a functional of the Cartan subalgebra of some Lie algebra (2). Indeed, for an integrable hamiltonian H we can find canonically conjugated action (I_a) and angle (θ^a) variables

$$\{I_a, \theta^b\}_\omega = \delta_a^b \quad (4)$$

such that the hamiltonian H is a functional of the action variables I_a only. These action variables label invariant torii on the phase space and the motion of $H(I)$ is constrained on these torii. Since the action variables are in involution,

$$\{I_a, I_b\}_\omega = 0 \quad (5)$$

we can interpret them as generators in the Cartan subalgebra of some Lie algebra (2).

As an example consider S^2 , which is the co-adjoint orbit of $SU(2)$. With z and $\bar{z}$ the coordinates on the Riemann sphere S^2 the Poisson bracket is

$$\{z, \bar{z}\} = \frac{1}{2iJ}(1 + |z|^2)^2 \quad (6)$$

This corresponds to the canonical spin- J representation of $SU(2)$ with generators

$$S_0 = -J \frac{1 - |z|^2}{1 + |z|^2} \quad S_+ = 2J \frac{\bar{z}}{1 + |z|^2} \quad S_- = 2J \frac{z}{1 + |z|^2} \quad (7)$$

Our hamiltonian $H(G_u)$ is then a generic functional of all generators (7), while an integrable hamiltonian is obtained *e.g.* by considering functions of S_0 only.

We are interested in the evaluation of the canonical phase space path integral

$$Z = \int [dz^a] \sqrt{||\omega_{ab}||} \exp\left\{i \int_0^\beta \vartheta_a \dot{z}^a - H(G_u)\right\} \quad (8)$$

for a generic classically integrable hamiltonian $H(G_u)$. Here ϑ_a are components of the symplectic one-form,

$$\omega_{ab} = \partial_a \vartheta_b - \partial_b \vartheta_a \quad (9)$$

In order to evaluate (8) we introduce arbitrary functions $j^u(t)$ that are independent of the phase space coordinates, and use familiar identities to write (8) as

$$Z = \exp\left\{-i \int_0^\beta H\left[\frac{1}{i} \frac{\delta}{\delta j^u(t)}\right]\right\} \int [dz^a] \sqrt{||\omega_{ab}||} \exp\left\{i \int_0^\beta \vartheta_a \dot{z}^a - j^u G_u\right\}_{|j=0} \quad (10)$$

Consequently it is sufficient to evaluate the following path integral

$$Z_o(j) = \int [dz^a] \sqrt{||\omega_{ab}||} \exp\left\{i \int_0^\beta \vartheta_a \dot{z}^a - j^u G_u\right\} \quad (11)$$

with arbitrary functions $j^u(t)$.

For j^u that are t -independent, (11) yields the Weyl character of the Lie group in the given representation. For arbitrary t -dependent functions $j^u(t)$ we consider an infinitesimal variation of the action in (11),

$$\delta(\vartheta_a \dot{z}^a - j^u G_u) = \delta z^a (\omega_{ab} \dot{z}^b - j^u \partial_a G_u) \quad (12)$$

We select

$$\delta z^a = \epsilon^u \{G_u, z^a\}_\omega \quad (13)$$

where the ϵ^u are some infinitesimal z^a -independent parameters. This corresponds to the infinitesimal limit of the canonical transformation

$$z^a \rightarrow e^{-\epsilon^u G_u} z^a e^{\epsilon^u G_u} = z^a + \epsilon^u \{z^a, G_u\} + \frac{1}{2} \epsilon^u \epsilon^v \{\{z^a, G_u\}, G_v\} + \dots \quad (14)$$

and yields

$$\delta(\vartheta_a \dot{z}^a - j^u G_u) = -\dot{\epsilon}^u G_u - f_{uv}{}^w \epsilon^u j^v G_w \quad (15)$$

Since the Liouville measure in (11) is invariant under canonical transformations we then conclude that the only effect of (13) and (15) in (11) is that the external sources $j^u(t)$ are shifted according to

$$j^u \rightarrow j^u + \dot{\epsilon}^u + f_{vw}{}^u \epsilon^v j^w \quad (16)$$

Notice that (16) is analogous to the nonabelian gauge transformation in a Yang-Mills theory. Indeed, if we relate the generators G_u to the Gauss law generators $G_u \sim D_i^{ab} E_i^b$ in a nonabelian gauge theory and identify $j^u(t)$ with the time component of the gauge field $j^u \sim A_o^a$, we observe that (16) has the functional form of a time dependent gauge transformation. On the basis of this analogy we then conclude that we can indeed expect the measure in (11) to be invariant under (13), and if it for some reason fails to be invariant we expect that the ensuing nontrivial Jacobian can be related to conventional nonabelian gauge anomalies.

As a consequence the path integral can only depend on gauge invariant functionals of j^μ *i.e.* the temporal Wilson loop. For the Cartan subalgebra T_u of G_u

$$\{T_u, T_v\} = 0 \quad (17)$$

corresponding to integrable models, we can then further simplify (16) to

$$j^u \rightarrow j^u + \dot{\epsilon}^u \quad (18)$$

and if we divide $j^u(t)$ into its constant part and its t -dependent part

$$j^u(t) = j_o^u + j_t^u \quad (19)$$

where j_o^u denotes the constant part and j_t^u the t -dependent part *e.g.* in a Fourier decomposition, we can gauge transform the path integral (11) to

$$Z_o(j) = \int [dz^a] \sqrt{\det ||\omega_{ab}||} \exp\{i \int \vartheta_a \dot{z}^a - j_o^u T_u\} \quad (20)$$

i.e. it depends *only* on the constant modes j_o^u . Since (10) is evaluated at $j^u(t) = 0$, we conclude that in the abelian case the original path integral (8) simplifies to

$$Z = \exp\left\{-i\beta H \left[\frac{1}{i} \frac{\partial}{\partial j_o^u}\right]\right\} \int [dz^a] \sqrt{\det|\omega_{ab}|} \exp\left\{i \int \vartheta_a \dot{z}^a - j_o^u T_u\right\}_{|j_o=0} \quad (21)$$

Consequently it is sufficient to evaluate the path integral

$$Z(j_o) = \int [dz^a][dc^a] \exp\left\{i \int \vartheta_a \dot{z}^a - T + \frac{1}{2} c^a \omega_{ab} c^b\right\} \quad (22)$$

where we have introduced anticommuting variables c^a to exponentiate the determinant of the symplectic two-form and

$$T = j_o^u T_u \quad (23)$$

is an element of the Cartan subalgebra *i.e.* a $U(1)$ generator of the Lie algebra,

The path integral in (22) yields the Weyl character of T and has been evaluated *e.g.* in [12,13] using localization methods. Here we need a few relevant steps: We interpret (22) as an integral in the loop space of the phase space, parametrized by the time evolution $z^a \rightarrow z^a(t)$. We identify $c^a(t)$ as a basis of loop space one-forms and interpret the bosonic part of the action in (22) as a loop space hamiltonian functional. The corresponding loop space hamiltonian vector field then has components

$$\mathcal{X}_S^a = \dot{z}^a - \omega^{ab} \partial_b T \quad (24)$$

We introduce the loop space exterior derivative (in the following time integrals are implicit)

$$d = c^a \partial_a \quad (25)$$

and a basis i_a of loop space contractions dual to the basis of one-forms $c^a(t)$,

$$i_a(t) c^b(t') = \delta_a^b(t - t') \quad (26)$$

We then define the following equivariant exterior derivative on the loop space,

$$d_S = d + i_S = c^a \partial_a + \mathcal{X}_S^a i_a \quad (27)$$

The corresponding Lie-derivative along the hamiltonian vector field (24) is

$$\mathcal{L}_S = d i_S + i_S d \quad (28)$$

On the subspace of the exterior algebra where (28) vanishes, (27) is then nilpotent.

The action in (22) is closed with respect to (27),

$$d_S(\vartheta_a \dot{z}^a - T + \frac{1}{2}c^a \omega_{ab} c^b) \quad (29)$$

In [10] it has been shown, that if we add to the action a d_S exact term,

$$S \rightarrow \int \vartheta_a \dot{z}^a - T + \frac{1}{2}c^a \omega_{ab} c^b + d_S \Psi \quad (30)$$

where Ψ is in the nilpotent subspace of (27),

$$\mathcal{L}_S \Psi = 0 \quad (31)$$

the corresponding path integral

$$Z_\Psi = \int [dz^a][dc^a] \exp\{i \int \vartheta_a \dot{z}^a - T + \frac{1}{2}c^a \omega_{ab} c^b + d_S \Psi\} \quad (32)$$

is independent of Ψ and coincides with (22). By selecting a suitable Ψ the path integral can then be evaluated by the localization method.

In order to construct the appropriate Ψ we observe that the phase space Γ admits a metric tensor g_{ab} which is Lie-derived by the $U(1)$ generator T ,

$$\mathcal{L}_T g = 0 \quad (33)$$

For this, it is sufficient to take any metric tensor on Γ and average it over the group. This yields a metric tensor that satisfies (33). If we then select

$$\Psi = \frac{1}{2}g_{ab} \dot{z}^a c^b \quad (34)$$

the condition (31) is satisfied and the corresponding action (32) is

$$S = \int \frac{1}{2}g_{ab} \dot{z}^a \dot{z}^b + (\vartheta_a - \frac{1}{2}g_{ab} \mathcal{X}_T^b) \dot{z}^a - T - \frac{1}{2}c^a (g_{ab} \partial_t + \dot{z}^c g_{bd} \Gamma_{ac}^d) c^b + \frac{1}{2}c^a \omega_{ab} c^b \quad (35)$$

Here Γ_{ac}^d is the Christoffel symbol for the metric tensor g_{ab} . We scale $g_{ab} \rightarrow \lambda g_{ab}$. According to the general argument [10] the path integral is independent of λ . Hence we can evaluate (32) in the $\lambda \rightarrow \infty$ and the result is [12,13]

$$Z_o = \int dz_o dc_o \exp\{-i\beta T + i\frac{\beta}{2}c_o^a \omega_{ab} c_o^b\} \frac{1}{\sqrt{\det[\delta_{ab}^d \partial_t - \frac{1}{2}(\Omega_{ab}^d + R_{ab}^d)]}} \quad (36)$$

where

$$\Omega_{ab}^d = \frac{1}{2}g^{ac}[\partial_b(g_{cd} \mathcal{X}_T^d) - \partial_c(g_{bd} \mathcal{X}_T^d)] \quad (37)$$

is the Riemannian momentum map of T [7] and $R^a_b = R^a_{bcd}c^c_o c^d_o$ is the Riemannian curvature two-form. If we evaluate the determinant using¹ *e.g.* ζ -function regularization we get

$$Z_o = \int dz_o dc_o \exp\{-i\beta T + i\frac{\beta}{2}c^a_o \omega_{ab} c^b_o\} \sqrt{\det \left[\frac{\frac{\beta}{2}(\Omega^a_b + R^a_b)}{\sinh[\frac{\beta}{2}(\Omega^a_b + R^a_b)]} \right]} \quad (38)$$

Here

$$Ch \left[\frac{\beta}{2}(T - \omega) \right] = \exp\{-i\beta T + i\frac{\beta}{2}c^a_o \omega_{ab} c^b_o\} \quad (39)$$

is the equivariant Chern character and

$$\hat{A} \left[\frac{\beta}{2}(\Omega^a_b + R^a_b) \right] = \sqrt{\det \left[\frac{\frac{\beta}{2}(\Omega^a_b + R^a_b)}{\sinh[\frac{\beta}{2}(\Omega^a_b + R^a_b)]} \right]} \quad (40)$$

is the equivariant $\hat{A}$ -genus. Both (39) and (40) are equivariantly closed,

$$(d + i_T)Ch \left[\frac{\beta}{2}(T - \omega) \right] = (d + i_T)\hat{A} \left[\frac{\beta}{2}(\Omega^a_b + R^a_b) \right] = 0 \quad (41)$$

If we substitute these into (21) we then obtain our integration formula for the path integral (8), (21)

$$Z = \exp\{-i\beta \cdot H \left[\frac{1}{i} \frac{\partial}{\partial j^u_o} \right]\} \cdot Ch \left[\frac{\beta}{2}(j^u_o T_u - \omega) \right] \hat{A} \left[\frac{\beta}{2}(j^u_o [\Omega_{T_u}]^a_b + R^a_b) \right]_{|j=0} \quad (42)$$

i.e. the path integral reduces to a derivative expansion of equivariant characteristic classes.

We shall now argue, that (42) is closely related to the (infinitesimal) Lefschetz number of a Dirac operator which is defined on the phase space Γ . For this we first consider the following Dirac operator on an even dimensional Riemannian manifold $\mathcal{M}$

$$\gamma^\mu D_\mu = \begin{pmatrix} 0 & D_+ \\ D_- & 0 \end{pmatrix} = \gamma^\mu (\partial_\mu + A_\mu + \frac{1}{8} \omega_{\mu ij} [\gamma^i, \gamma^j]) \quad (43)$$

Here $\gamma^i = e^i_\mu \gamma^\mu$ are local Dirac matrices, e^i_μ are components of the vielbein and $\omega_{\mu ij}$ are components of the spin connection. We shall assume that the manifold $\mathcal{M}$ admits an operator realization of the Lie algebra (2),

$$[\mathcal{G}_u, \mathcal{G}_v] = f_{uv}^w \mathcal{G}_w \quad (44)$$

¹ Here we use a regularization which is symmetric in the negative and positive eigenvalues of the differential operator in (36)

such that the operators $\mathcal{G}_u$ commute with the Dirac operator (43)

$$[\mathcal{G}_u, \gamma^\mu D_\mu] = 0 \quad (45)$$

The eigenstates of $\gamma^\mu D_\mu$ that correspond to a fixed eigenvalue λ then define a representation of the Lie algebra $\mathcal{G}_u$. We shall be particularly interested in the representation which is defined by the zeromodes $\lambda = 0$ of $\gamma^\mu D_\mu$. For this we introduce some arbitrary parameters α_u and construct the linear combination $\alpha \cdot \mathcal{G}$ which is an element of the Cartan subalgebra of (44) in a suitable basis. We then introduce the following Lefschetz number,

$$L(\alpha \cdot \mathcal{G}) = \lim_{\beta \rightarrow \infty} \text{Tr} \{ e^{i\alpha \cdot \mathcal{G}} (e^{-\beta D_+ D_-} - e^{-\beta D_- D_+}) \} \quad (46)$$

One can show that the trace on the *r.h.s.* of (46) is independent of β and provided either $D_+ D_-$ or $D_- D_+$ has no zero modes it coincides with the character of the (reducible or irreducible) representation of $\mathcal{G}_u$ determined by the zeromodes of (43).

We wish to relate the evaluation of the Lefschetz number (46) to the evaluation of the path integral (8). For this we need the appropriate canonical realization of the Dirac operator $\gamma^\mu D_\mu$ and the Lie algebra generators $\mathcal{G}_u$: We interpret the coordinates z^μ on $\mathcal{M}$ as canonical coordinates on a phase space with conjugate momenta p_μ and Poisson brackets

$$\{p_\mu, z^\nu\} = \delta_\mu^\nu \quad (47)$$

We also realize the local γ -matrix algebra canonically using anticommuting variables ψ^i with graded Poisson brackets

$$\{\psi^i, \psi^j\} = \eta^{ij} \quad (48)$$

where η^{ij} is the local flat metric. The corresponding $\mathcal{M}$ -dependent variables $\psi^\mu = E_i^\mu \psi^i$ have the brackets

$$\{\psi^\mu, \psi^\nu\} = g^{\mu\nu} \quad (49)$$

where $g_{\mu\nu}$ is the metric tensor on $\mathcal{M}$. The appropriate canonical realization of the Dirac operator (43) is then

$$\mathcal{Q} = \psi^\mu \mathcal{P}_\mu = \psi^\mu (p_\mu + \frac{1}{4} \omega_{\mu ij} \psi^i \psi^j + A_\mu) \quad (50)$$

and

$$\{\mathcal{Q}, \mathcal{Q}\} = \mathcal{H} = g_{\mu\nu} \mathcal{P}^\mu \mathcal{P}^\nu + \frac{1}{2} \psi^\mu \psi^\nu F_{\mu\nu} =$$

$$= g^{\mu\nu}(p_\mu + \frac{1}{4}\omega_{\mu ij}\psi^i\psi^j + A_\mu)(p_\nu + \frac{1}{4}\omega_{\mu kl}\psi^k\psi^l + A_\nu) + \frac{1}{2}\psi^\mu\psi^\nu F_{\mu\nu} \quad (51)$$

where we have used identities of the Riemann tensor.

We first consider the path integral for the Atiyah-Singer index of (43), *i.e.* we set $\alpha = 0$ in (46). This path integral is (see *e.g.* [11])

$$\int [dz^\mu][dp_\mu][d\psi^\mu] \exp\{i \int p_\mu \dot{z}^\mu - \frac{1}{2}\psi_\mu \dot{\psi}^\mu - g_{\mu\nu} \mathcal{P}^\mu \mathcal{P}^\nu - \frac{1}{2}\psi^\mu\psi^\nu F_{\mu\nu}\} \quad (52)$$

We change variables

$$p_\mu \rightarrow p_\mu - A_\mu - \omega_{\mu ij}\psi^i\psi^j \quad (53)$$

This gives

$$S = \int p_\mu \dot{z}^\mu - A_\mu \dot{z}^\mu - \frac{1}{2}g_{\mu\nu} p^\mu p^\nu - \frac{1}{2}\psi^\mu (g_{\mu\nu} \partial_t + \dot{z}^\rho g_{\mu\sigma} \Gamma_{\rho\nu}^\sigma) \psi^\nu - \frac{1}{2}\psi^\mu \omega_{\mu\nu} \psi^\nu \quad (54)$$

and eliminating p_μ we get

$$S = \int \frac{1}{2}g_{\mu\nu} \dot{z}^\mu \dot{z}^\nu - A_\mu \dot{z}^\mu - \frac{1}{2}\psi^\mu (g_{\mu\nu} \partial_t + \dot{z}^\rho g_{\mu\sigma} \Gamma_{\rho\nu}^\sigma) \psi^\nu - \frac{1}{2}\psi^\mu F_{\mu\nu} \psi^\nu \quad (55)$$

With the identification $A_\mu \sim -\vartheta_a$ and $F_{\mu\nu} \sim -\omega_{ab}$ this coincides with the action in (35) with $T = 0$.

We shall now proceed to demonstrate that if $\alpha \neq 0$ in (46), its path integral representation coincides with (35) for an appropriate $T \neq 0$. For this we need an operator realization of the Lie algebra (44) in the extended phase space. If we define

$$\mathcal{G}_u = \mathcal{X}_u^\mu \frac{\partial}{\partial z^\mu} - \mathcal{X}_{u,\nu}^\mu p^\nu \frac{\partial}{\partial p^\mu} + \mathcal{X}_{u,\nu}^\mu \psi^\nu \frac{\partial}{\partial \psi^\mu} \quad (56)$$

where $\mathcal{X}_u^\mu(z)$ are components of the hamiltonian vector fields corresponding to the functions $G_u(z)$ in (2), we find that (56) satisfy the Lie algebra (44).

From (54) we extract the following symplectic one-form on the extended phase space,

$$\Theta = (p_\mu + \vartheta_\mu - \frac{1}{2}\psi^\nu g_{\nu\sigma} \Gamma_{\mu\rho}^\sigma \psi^\rho) dz^\mu - \frac{1}{2}\psi^\mu g_{\mu\nu} d\psi^\nu \quad (57)$$

We shall assume that in the original phase space the symplectic one-form $\vartheta_\mu(z) dz^\mu$ is Lie derived by the vector fields $\mathcal{X}_u^a$,

$$\mathcal{L}_{G_u} \vartheta = 0 \quad (58)$$

As a consequence we find that in the extended phase space (56) Lie derives (57),

$$\mathcal{L}_{\mathcal{G}_u} \Theta = 0 \quad (59)$$

Integrating this we then obtain the momentum map

$$i_{\mathcal{G}_u}\Theta = \mathcal{T}_u = \mathcal{X}_u^\mu p_\mu + \vartheta_\mu \mathcal{X}_u^\mu - \frac{1}{2}\psi^\mu \Omega_{\mu\nu}^u \psi^\nu \quad (60)$$

where

$$\Omega_{\mu\nu}^u = \frac{1}{2}[\partial_\mu(g_{\nu\sigma}\mathcal{X}_u^\sigma) - \partial_\nu(g_{\mu\sigma}\mathcal{X}_u^\sigma)] \quad (61)$$

is the Riemannian momentum map for the generator G_u . In the extended phase space equipped with the symplectic structure defined by (57) the momentum maps (60) determine a Poisson bracket realization of (2),

$$\{\mathcal{T}_u, \mathcal{T}_v\} = f_{uv}{}^w \mathcal{T}_w \quad (62)$$

Furthermore, since

$$\{\mathcal{T}_u, \mathcal{Q}\} = 0 \quad (63)$$

we conclude that the path integral for the action

$$S = \int p_\mu \dot{z}^\mu + \vartheta_\mu \dot{z}^\mu - \frac{1}{2}\psi^\mu(g_{\mu\nu}\partial_t + g_{\mu\sigma}\dot{z}^\rho \Gamma_{\rho\nu}^\sigma)\psi^\nu + \frac{1}{2}\omega_{\mu\nu}\psi^\mu\psi^\nu - \frac{1}{2}g_{\mu\nu}p^\mu p^\nu - \tilde{\alpha} \cdot \mathcal{T} \quad (64)$$

where $\tilde{\alpha} = \beta^{-1}\alpha$, yields the Lefschetz number (46). Eliminating p_μ we get further

$$\begin{aligned} S = \int & \frac{1}{2}g_{\mu\nu}\dot{z}^\mu\dot{z}^\nu + \dot{z}^\mu[\vartheta_\mu + g_{\mu\nu}(\tilde{\alpha} \cdot \mathcal{X}^\mu)] + \frac{1}{2}g_{\mu\nu}(\tilde{\alpha} \cdot \mathcal{X}^\mu)(\tilde{\alpha} \cdot \mathcal{X}^\nu) + (\tilde{\alpha} \cdot \mathcal{X}^\mu)\vartheta_\mu - \\ & - \frac{1}{2}\psi^\mu(g_{\mu\nu}\partial_t + \dot{z}^\rho g_{\mu\sigma}\Gamma_{\rho\nu}^\sigma)\psi^\nu + \frac{1}{2}\psi^\mu\omega_{\mu\nu}\psi^\nu + \frac{1}{2}\psi^\mu(\tilde{\alpha} \cdot \Omega_{\mu\nu})\psi^\nu \end{aligned} \quad (65)$$

If we select ϑ_μ so that T is obtained as the momentum map

$$T = -(\tilde{\alpha} \cdot \mathcal{X}^\mu)\vartheta_\mu \quad (66)$$

we then find

$$\begin{aligned} S = \int & \frac{1}{2}g_{\mu\nu}\dot{z}^\mu\dot{z}^\nu + \dot{z}^\mu(\vartheta_\mu + g_{\mu\nu}\mathcal{X}_T^\mu) + \frac{1}{2}g_{\mu\nu}\mathcal{X}_T^\mu\mathcal{X}_T^\nu - T - \\ & - \frac{1}{2}\psi^\mu(g_{\mu\nu}\partial_t + \dot{z}^\rho g_{\mu\sigma}\Gamma_{\rho\nu}^\sigma)\psi^\nu + \frac{1}{2}\psi^\mu\omega_{\mu\nu}\psi^\nu + \frac{1}{2}\psi^\mu\Omega_{\mu\nu}^T\psi^\nu \end{aligned} \quad (67)$$

This action has the following interpretation: From (55) we conclude that the action

$$S_1 = \int \frac{1}{2}g_{\mu\nu}\dot{z}^\mu\dot{z}^\nu - \frac{1}{2}\psi^\mu(g_{\mu\nu}\partial_t + \dot{z}^\rho g_{\mu\sigma}\Gamma_{\rho\nu}^\sigma)\psi^\nu \quad (68)$$

can be interpreted as a Dirac operator on the Riemannian manifold. Similarly, the path integral for

$$S_2 = \int \vartheta_\mu \dot{z}^\mu - T + \frac{1}{2}\psi^\mu\omega_{\mu\nu}\psi^\nu \quad (69)$$

yields the Weyl character of T . Finally, the action

$$S_3 = \int g_{\mu\nu} \mathcal{X}_T^\mu \dot{z}^\mu - \frac{1}{2} g_{\mu\nu} \mathcal{X}_T^\mu \mathcal{X}_T^\nu + \frac{1}{2} \psi^\mu \Omega_{\mu\nu}^T \psi^\nu \quad (70)$$

defines a bi-hamiltonian pair of S_2 in the sense that in the place of the symplectic structure (9) we have

$$\partial_\mu \left(\frac{1}{2} g_{\nu\rho} \mathcal{X}_T^\rho \right) - \partial_\nu \left(\frac{1}{2} g_{\mu\rho} \mathcal{X}_T^\rho \right) = \Omega_{\mu\nu}^T \quad (71)$$

and in the place of the hamiltonian T we have

$$\mathcal{T} = \frac{1}{2} \psi^\mu \Omega_{\mu\nu}^T \psi^\nu \quad (72)$$

and the classical equations of motion for the pairs (T, ω) and $(\mathcal{T}, \Omega)$ coincide,

$$\dot{z}^\mu = \{T, z^\mu\}_\omega = \{\mathcal{T}, z^\mu\}_\Omega \quad (73)$$

The existence of this bi-hamiltonian structure is of course consistent with the classical integrability of the hamiltonian system (T, ω) .

In order to relate the actions (35) and (67) we introduce

$$\tilde{\Psi} = \frac{1}{2} g_{\mu\nu} \mathcal{X}_T^\mu \psi^\nu \quad (74)$$

and add the corresponding d_S -exact term to the action (35),

$$S \rightarrow S + d_S \tilde{\Psi} \quad (75)$$

This action then coincides with (67), and provided either $D_+ D_-$ or $D_- D_+$ in (46) has no zeromodes, the corresponding path integrals must also coincide (possibly modulo an overall sign). Furthermore, from (22) and (32) we conclude that the path integral for (67) differs from the path integral for the Weyl character only by a d_S exact term. Consequently the infinitesimal Lefschetz number (46) of our Dirac operator coincides with the Weyl character of T , provided either $D_+ D_-$ or $D_- D_+$ in (46) has no zeromodes and the boundary conditions for the path integrals have been properly selected.

In conclusion, we have generalized the localization evaluation of path integrals discussed in [12,13] to a general class of hamiltonians which are constructed from Lie algebra generators. We have also established that the path integral for these hamiltonians evaluates to equivariant characteristic classes, and is also closely related to the Lefschetz number of a Dirac operator. Our results indicate that equivariant cohomology could provide a natural geometric framework for understanding quantum integrability.

We thank A. Alekseev, L. Faddeev, V. Fock, A. Gerasimov, N. Nekrasov, A. Rosly and O. Tirkkonen for discussions.

References

1. L.D. Faddeev and L.A. Takhtajan, *Hamiltonian Methods in the Theory of Solitons* (Springer Verlag 1987)
2. A. Das, *Integrable Models* (World Scientific 1989)
3. J.J. Duistermaat and G.J. Heckman, *Inv. Math.* **69** (1982) 259; and *ibid* **72** (1983) 153
4. N. Berline and M. Vergne, *Duke Math. Journ.* **50** (1983) 539
5. M.F. Atiyah and R. Bott, *Topology* **23** (1984) 1
6. J.-M. Bismut, *Comm. Math. Phys.* **98** (1985) 213; and *ibid.* **103** (1986) 127; and *J. Funct. Anal.* **62** (1985) 435
7. N. Berline, E. Getzler and M. Vergne, *Heat Kernels and Dirac Operators* (Springer verlag, Berlin, 1991)
8. M.F. Atiyah, *Asterisque* **131** (1985) 43
9. E. Witten, *J. Geom. Phys.* **9** (1992) 303
10. M. Blau, E. Keski-Vakkuri and A.J. Niemi, *Phys. Lett.* **B246** (1990) 92;
11. A. Hietamäki, A.Yu. Morozov, A.J. Niemi and K. Palo, *Phys. Lett.* **263B** (1991) 417; A. Yu. Morozov, A.J. Niemi and K. Palo, *Phys. Lett.* **271** (1991) 365; and *Nucl. Phys.* **B377** (1992) 295
12. A.J. Niemi and O. Tirkkonen, *Phys. Lett.* **B293** (1992) 339
13. A.J. Niemi and O. Tirkkonen, preprint hep-th-9301059